

ADDITIVE K -COLORABLE EXTENSIONS OF THE RATIONAL PLANE

Klaus G. FISCHER

Department of Mathematical Sciences, George Mason University, Fairfax, VA 22030, USA

Received 4 December 1987

Revised 24 May 1988

Let F be a field, $\mathbb{Q} \subset F \subset \mathbb{R}$ and consider F^d as a graph with vertices the points of F^d and an edge between two points if their Euclidean distance is 1. Let $\Sigma_0(F^d)$ be the subgroup of F^d generated by the unit vectors ξ . If G is a group of order k , then a group homomorphism $v: \Sigma_0(F^d) \rightarrow G$ for which $v(\xi) \neq 0$ whenever $\|\xi\| = 1$ is said to be an additive k -coloring of F^d . The known 2 and 4-colorings of \mathbb{Q}^3 and \mathbb{Q}^4 respectively, are shown to be additive. If N is a square free integer, then it is shown that $\mathbb{Q}(\sqrt{N})^2$ has an additive 2-coloring iff $N \not\equiv 3 \pmod{4}$. If $N \not\equiv 2 \pmod{3}$, then $\mathbb{Q}(\sqrt{N})^2$ has an additive 3-coloring. Hence, it follows that the chromatic number of $\mathbb{Q}(\sqrt{3})^2$ is 3. The existence of additive colorings on $\mathbb{Q}(\sqrt{N})^2$ for the remaining cases is also discussed.

Additive k -colorings constrain cycles in F^d to satisfy group identities. Hence, it is shown for example, that if F^2 is 2-colorable and if $\sqrt{2} \notin F$, then F^2 contains no regular polygon except for the square. This generalizes the classical result known for the rational plane.

1. Introduction

Let F be a subfield of the reals \mathbb{R} so that $\mathbb{Q} \subset F \subset \mathbb{R}$ where \mathbb{Q} is the field of rationals. By F^d is meant the collection of points in the real Euclidean space \mathbb{R}^d all of whose d coordinates are in F . These points will be interpreted as vertices of a graph, also denoted by F^d , with an edge between the vertices P and Q if their Euclidean distance $\|P - Q\| = 1$. The connected component containing the origin will be denoted by $\Sigma_0(F^d)$. Hence, this is the subgroup in F^d generated by all the unit vectors.

If k is a positive integer, a k -coloring of F^d is an assignment v of the vertices to a set of k elements so that $v(P) \neq v(Q)$ whenever $\|P - Q\| = 1$. It is easy to see that F^d can be k -colored iff $\Sigma_0(F^d)$ can be k -colored. A group homomorphism $v: \Sigma_0(F^d) \rightarrow G$ to an additive group G of order k so that $v(\xi) \neq 0$ whenever $\|\xi\| = 1$, is said to be an additive k -coloring of F^d .

An additive k -coloring clearly provides a k -coloring of F^d . The converse of this is not true although it easy to see that F^d can be 2-colored iff it possesses an additive 2-coloring. It is shown, for example, that \mathbb{Q}^2 , \mathbb{Q}^3 and \mathbb{Q}^4 have additive 2, 2 and 4-colorings respectively. That these are also the chromatic numbers of the spaces was known to Woodall [9], Johnson [7] and to Benda and Perles [3] respectively. Quite generally it is shown that if α is a real transcendental over F , then F^d has an additive k -coloring iff $F(\alpha)^d$ does. The same statement holds true

for non-additive k -colorings. Additive k -colorings place restrictions on the cycles in F^d since these must now satisfy group identities. Thus, for example, if F^2 is two colorable and $\sqrt{2} \notin F$, then F^2 contains no regular polygon except for the square. In view of the fact that \mathbb{Q}^2 is two-colorable, this generalizes the known result for the rational plane [6, p. 4].

This paper considers possible additive k -colorings of the plane formed by a real quadratic extension of \mathbb{Q} . If N is a square free positive integer, it is shown that the plane $\mathbb{Q}(\sqrt{N})^2$ has an additive 2-coloring iff $N \not\equiv 3 \pmod{4}$. That $\mathbb{Q}(\sqrt{N})^2$ has a two coloring in this case also follows by a recent result of Johnson's in [8]. The converse answers a problem posed by him in that paper. If $N \not\equiv 2 \pmod{3}$, it is shown here that $\mathbb{Q}(\sqrt{N})^2$ has an additive 3-coloring. It follows, for example, that the chromatic number of $\mathbb{Q}(\sqrt{3})^2$ is 3. Furthermore, if $N \equiv 3 \pmod{8}$, then $\mathbb{Q}(\sqrt{N})^2$ has an additive 4-coloring and therefore places 4 as a bound on its chromatic number. In the remaining cases, it is shown that $1/k \in \Sigma_0(\mathbb{Q}(\sqrt{N})^2)$ for $1 \leq k \leq 6$. Hence, no additive k -coloring for these values of k can exist since if $v: \Sigma_0(\mathbb{Q}(\sqrt{N})^2) \rightarrow G$ is such a coloring, then $k \cdot v(1/k) = v(1) \neq 0$ and yet $k \cdot G = 0$. In fact, $\mathbb{Q}(\sqrt{167})^2$ does not allow an additive k -coloring for $k < 11$.

In view of the fact that the chromatic number of \mathbb{R}^2 is at least 4 but not more than 7 [5, p. 16], the above example among others shows that additive k -colorings even on quadratic extensions of \mathbb{Q} are too special to provide k -colorings in most cases. This still begs the general question, however, whether some algebraic criterion can be found to determine the chromatic number of F^2 where F is a subfield of \mathbb{R} , of finite degree over \mathbb{Q} . That the answer to this question would suffice to find the chromatic number of \mathbb{R}^2 has already been noted by Benda and Perles in [2]. The assertion also follows easily here from the result of de Bruijn and Erdős in [1] which states that the chromatic number of \mathbb{R}^2 is the same as the number of colors required to paint any finite number of points in \mathbb{R}^2 , and the above theorem on adjoining transcendentals.

2. Additive k -colorings

Let $\Sigma_0(F^d)$ be the points P in the graph F^d for which there exists a path in F^d from the origin 0 to P . It should be emphasized that this means there exists a sequence of points $0 = P_0, \dots, P_n = P$ in F^d so that $\|P_j - P_{j-1}\| = 1$ for $j = 1, \dots, n$. Clearly,

$$\Sigma_0(F^d) = \left\{ \xi : \xi = \sum_{j=1}^n \xi_j, \xi_j \in F^d \text{ and } \|\xi_j\| = 1 \text{ for all } j \right\}$$

and it forms an additive subgroup of F^d . The translates $\Sigma_Q(F^d) = Q + \Sigma_0(F^d)$ identify the points P in F^d which may be reached by a path in F^d from the point Q in F^d .

Since it is always assumed that the field $F \subset \mathbb{R}$, the plane F^2 may be identified with the field of complex numbers $F(i)$. Hence, $\Sigma_0(F^2)$ may be identified as sums

of complex numbers in $F(i)$ of norm 1 which form a ring containing the integers \mathbb{Z} .

In what follows, any group G is considered to be abelian and additive with identity element 0. The ring $\mathbb{Z}/(2)$ will denote the ring of integers modulo 2 and $[n]$ the equivalence class of an integer n modulo 2.

Theorem 1. i. If F^d has an additive k -coloring, then F^d can be k -colored.

ii. F^d can be 2-colored iff there exists an additive map $v: \Sigma_0(F^d) \rightarrow \mathbb{Z}/(2)$ so that $v(\xi) = [1]$ if $\|\xi\| = 1$.

iii. The plane F^2 can be 2-colored iff there exists a non-zero ring homomorphism $v: \Sigma_0(F^2) \rightarrow \mathbb{Z}/(2)$.

Proof. Since the connected components of F^d are translates of $\Sigma_0(F^d)$, to k -color F^d it suffices to k -color $\Sigma_0(F^d)$. The existence of a group homomorphism $v: \Sigma_0(F^d) \rightarrow G$ where G is of order k , assures that if $\xi, \eta \in \Sigma_0(F^d)$ and if $\|\xi - \eta\| = 1$, then $v(\xi - \eta) \neq 0$. Hence, $v(\xi) \neq v(\eta)$ and therefore v assigns a k -coloring to $\Sigma_0(F^d)$. This provides the proof for part (i).

To show (ii), note that if F^d can be 2-colored, then the sequence of points in a path must have alternating colors and hence every cycle must have an even number of edges. It follows that if $\xi = \sum_{j=1}^n \xi_j \in \Sigma_0(F^d)$, with $\|\xi_j\| = 1$, the map $v: \Sigma_0(F^d) \rightarrow \mathbb{Z}/(2)$ for which $v(\xi) = [n]$ is well defined. It is easily seen to be additive and $v(\xi) = [1]$ if $\|\xi\| = 1$. Using part (i), this proves (ii).

For a 2-colorable plane F^2 , the additive map from (ii) is easily seen to be a ring homomorphism since by assumption $v(\xi_1 \cdot \xi_2) = [1]$ if $\|\xi_1\| = \|\xi_2\| = 1$. But conversely, given any non-zero homomorphism on the ring $\Sigma_0(F^2)$, it is clearly additive and if $\|\xi\| = 1$, then ξ has an inverse in $\Sigma_0(F^2)$ and therefore $v(\xi) = 1$. It follows from (i) that F^2 is 2-colorable and proves part (iii). \square

Let $\mathbb{Z}_{(2)}$ be the ring of fractions of \mathbb{Q} with odd denominators. This is a local ring with maximal ideal (2) . Let $\epsilon: \mathbb{Z}_{(2)} \rightarrow \mathbb{Z}/(2)$ be the evaluation at the maximal ideal.

Note that for any integer n , $n \equiv 0 \pmod{2}$ iff $n^2 \equiv 0 \pmod{4}$; and $n \equiv 1 \pmod{2}$ iff $n^2 \equiv 1 \pmod{4}$. As an immediate application of the theorem one has the following.

Theorem 2. \mathbb{Q}^3 has an additive 2-coloring.

Proof. If $\xi = (a/d, b/d, c/d) \in \mathbb{Q}^3$ with $\|\xi\| = 1$, then it may be assumed that the integers satisfy $(a, b, c, d) = 1$ and that

$$a^2 + b^2 + c^2 = d^2.$$

Considering the equation $a^2 + b^2 + c^2 = d^2 \pmod{4}$, it can be assumed that $d \equiv 1 \pmod{2}$ since $(a, b, c, d) = 1$. Hence, $\Sigma_0(\mathbb{Q}^3) \subset \mathbb{Z}_{(2)}^3$. Furthermore, it follows that exactly one of the integers a, b , or c must equal $1 \pmod{2}$ and hence $[a + b + c] = [1]$.

Let $s: \mathbb{Q}^3 \rightarrow \mathbb{Q}$, where $s(a, b, c) = a + b + c$, be the additive linear functional which sums the coordinates of \mathbb{Q}^3 . Since $\Sigma_0(\mathbb{Q}^3) \subset \mathbb{Z}_{(2)}^3$ by the above, the restriction of s together with the map ϵ gives a mapping of groups

$$v = \epsilon \cdot s: \Sigma_0(\mathbb{Q}^3) \rightarrow \mathbb{Z}/(2),$$

for which if $\|\xi\| = 1$, then $v(\xi) = [a + b + c] = [1]$. \square

Although \mathbb{Q}^3 may be 2-colored it does allow equilateral triangles. (But not with sides of rational length, of course, since a dilation would contradict the above theorem.) For example, the points

$$(1, 0, 0), (0, 1, 0) \text{ and } (0, 0, 1)$$

give an equilateral triangle for which the length of each side is the irrational number $\sqrt{2}$. That this, in particular, is not the case for a 2-colorable plane F^2 is a consequence of the following theorem.

Theorem 3. *Let F be a field contained in \mathbb{R} and let $r \neq 0$ be a fixed real number that may be realized as the distance between two points of F^2 . Let P_0, \dots, P_n with $P_0 = P_n$ be a sequence of points in the plane F^2 so that $\|P_j - P_{j-1}\| = r$ for each $j = 1, \dots, n$. Then F^2 is 2-colorable iff $n = 0 \pmod{2}$ for any such sequence of points.*

Proof. Assume that there exists $\eta_j \in F(i)$ for $j = 1, \dots, n$ with $\|\eta_j\| = r$ and

$$0 = \eta_1 + \dots + \eta_n.$$

Since η_1^{-1} is also in $F(i)$, it follows that

$$0 = \eta_1^{-1} \cdot \eta_1 + \dots + \eta_1^{-1} \cdot \eta_n \in \Sigma_0(F^2),$$

since $\|\eta_1^{-1} \cdot \eta_j\| = 1/r \cdot r = 1$ for each j . If F is 2-colorable, applying the map v from Theorem 1 shows that $n = 0 \pmod{2}$.

Conversely, to say that $r \neq 0$ is realized as the distance between two points of F^2 says there exist some nonzero $\eta \in F^2$ for which $\|\eta\| = r$. If $\sum_{j=1}^n \xi_j = 0$ where $\|\xi_j\| = 1$, then $\sum_{j=1}^n \eta \cdot \xi_j = 0$ yields a sequence of points P_0, \dots, P_n in F^2 with $P_0 = P_n$ and $\|P_j - P_{j-1}\| = r$ for each j . The assumption on such a sequence assures that $n = 0 \pmod{2}$. As in Theorem 1, it follows that for any $\sum_{j=1}^n \xi_j \in \Sigma_0(F^2)$, the assignment $v(\sum_{j=1}^n \xi_j) = [n]$ is well defined and provides an additive 2-coloring. Therefore, F^2 is 2-colorable. \square

Note that the theorem demonstrates the fact that no suitable multiplication of vectors in \mathbb{Q}^3 can be found to be compatible with the Euclidean norm. For if this were so, then since \mathbb{Q}^3 is 2-colorable, the same proof would contradict the existence of equilateral triangles in \mathbb{Q}^3 .

The next theorem shows that regular n -gons for $n \neq 4$ cannot exist in certain 2-colorable planes.

Theorem 4. *Let F be a field contained in \mathbb{R} for which the plane F^2 is 2-colorable and assume that $\sqrt{2} \notin F$. Then the only regular polygon in the plane F^2 is the square.*

Proof. To say that a regular n -polygon exists in the plane F^2 means we can find a primitive n th root of unity $\xi \in F(i)$ with $n \geq 3$. Hence $\xi^n = 1$ and

$$0 = 1 + \xi + \cdots + \xi^{n-1}.$$

If p is a prime divisor of n for which $n = p \cdot m$ for some positive integer m , then

$$\begin{aligned} 0 &= (1 + \xi + \cdots + \xi^{m-1}) + (1 + \xi + \cdots + \xi^{m-1}) \cdot \xi^m + \cdots \\ &\quad + (1 + \xi + \cdots + \xi^{m-1}) \cdot (\xi^m)^{p-1}. \end{aligned}$$

Since $(1 + \xi + \cdots + \xi^{m-1}) \neq 0$, it follows that

$$1 + \xi^m + \cdots + (\xi^m)^{p-1} = 0.$$

Since $\|\xi^j\| = 1$ for all integers j and since F^2 is 2-colorable, it follows that p must be even and hence $n = 2^s$ for some positive integer s . But if $s \geq 3$, then the complex number $\eta = \xi^{2^{s-3}}$ is a primitive 8th root of unity. Hence, $\eta = (1/\sqrt{2}) \times (\pm 1 + \pm i) \in F(i)$. But this cannot be since $\sqrt{2} \notin F$. Hence $s = 2$ and therefore ξ must be a 4th root of unity. This root corresponds to the square which certainly exists in F^2 . \square

The theorem gives as a consequence a very geometrical proof of the following classical result [6, p. 4].

Corollary. *The only regular polygon in the rational plane is the square.*

Proof. Since \mathbb{Q}^2 is 2-colorable by Theorem 2 and $\sqrt{2} \notin \mathbb{Q}$, the result follows from the theorem. \square

The explicit description of the points P in $\Sigma_0(F^d)$ is of interest in its own right since these are the ones that may be reached by a path in F^d from the origin. Of course $\Sigma_0(\mathbb{Q}^1) = \mathbb{Z}$ and using Lagrange's Four Square Theorem, it is easily seen in [3] or [2] that $\Sigma_0(\mathbb{Q}^d) = \mathbb{Q}^d$ for $d \geq 5$. In fact, Chilakamarri in [3] computes the maximum number of points in \mathbb{Q}^d that are pairwise a unit distance apart. This is the clique number of the graph whose vertices are points in \mathbb{Q}^d with vertices adjacent if their distance is 1.

Let $S \subset \mathbb{Z}$ be the multiplicative set given by $S = \{p_1^{\alpha_1} \cdots p_r^{\alpha_r} : \alpha_j \text{ nonnegative integers and } p_j \text{ are primes for which } p_j \equiv 1 \pmod{4} \forall j\}$ and let $\mathbb{Z}_S = \{m/s : m \in \mathbb{Z}, s \in S\}$ denote the localization of \mathbb{Z} at S . Recorded below is the description of $\Sigma_0(\mathbb{Q}^d)$ for $d = 2, 3$, and 4. It will be seen from these that \mathbb{Q}^4 has an additive 4 coloring.

Example 1. $\Sigma_0(\mathbb{Q}^2) = \mathbb{Z}_S + \mathbb{Z}_S \cdot i$ [2].

If $\xi = a/c + i \cdot b/c \in \Sigma_0(\mathbb{Q}^2)$ with $\|\xi\| = 1$, then it may be assumed that a , b and c form a primitive Pythagorean triplet. That is,

$$a^2 + b^2 = c^2 \quad \text{and} \quad (a, b, c) = 1.$$

Since $(a, b, c) = 1$, the last equation taken modulo p shows that no prime $p = 3 \pmod{4}$ can occur as a factor of the denominator c of such a triplet. Neither can the prime 2. Hence, $\Sigma_0(\mathbb{Q}^2) \subset \mathbb{Z}_S + \mathbb{Z}_S \cdot i$.

To show the opposite inclusion, it suffices to show that $\mathbb{Z}_S \subset \Sigma_0(\mathbb{Q}^2)$. Since $\Sigma_0(\mathbb{Q}^2)$ is a ring, and since $\mathbb{Z} \subset \Sigma_0(\mathbb{Q}^2)$, it therefore suffices to show that for a prime $p = 1 \pmod{4}$, $1/p \in \mathbb{Z}_S$. For such a p we have integers a and b with $(a, p) = (b, p) = 1$ and $p^2 = a^2 + b^2$. Therefore, $\xi = a/p + b/p \cdot i$ has norm 1. But then if $\bar{\xi}$ is the complex conjugate of ξ , it follows that $\bar{\xi} + \xi = (2 \cdot a)/p \in \Sigma_0(\mathbb{Q}^2)$. Since $(p, 2a) = 1$, we may find integers m , n so that $m \cdot p + n \cdot 2 \cdot a = 1$. It follows that

$$\frac{1}{p} = \frac{m \cdot p + n \cdot 2 \cdot a}{p} = m + \frac{n \cdot 2 \cdot a}{p} \in \Sigma_0(\mathbb{Q}^2),$$

since both m and $(n \cdot 2 \cdot a)/p$ are in the ring $\Sigma_0(\mathbb{Q}^2)$.

Example 2. $\Sigma_0(\mathbb{Q}^3) = \mathbb{Z}_{(2)}^3$.

If $(a/d, b/d, c/d)$ is a unit vector where a , b , c and d are integers for which $(a, b, c, d) = 1$, the equation $a^2 + b^2 + c^2 = d^2$ taken modulo 4 shows that $[d] = [1]$. Hence any point in $\Sigma_0(\mathbb{Q}^3)$ must have an odd denominator.

To show that $\Sigma_0(\mathbb{Q}^3) \supset \mathbb{Z}_{(2)}^3$, it suffices to show that $(1/d, 0, 0) \in \Sigma_0(\mathbb{Q}^3)$ where d is odd. But if $d = e \cdot f$ where $(e, f) = 1$, then $1/e + 1/f = (f + e)/d$ where $(d, f + e) = 1$. As in the above example, one shows that if $(a/d, 0, 0) \in \Sigma_0(\mathbb{Q}^3)$ where $(a, d) = 1$, then $(1/d, 0, 0) \in \Sigma_0(\mathbb{Q}^3)$. Hence, it is sufficient to show that $(1/d, 0, 0) \in \Sigma_0(\mathbb{Q}^3)$ where $d = p^k$ is a positive power k of an odd prime p .

By the previous example, if $p = 1 \pmod{4}$ then $(1/d, 0) \in \Sigma_0(\mathbb{Q}^2)$ and therefore $(1/d, 0, 0) \in \Sigma_0(\mathbb{Q}^3)$.

For the case that $p = 3 \pmod{4}$, a result credited to E. Lionnet [4] states that any odd integer may be written as the sum of four squares two of which are consecutive. Hence d may be written as $d = x^2 + y^2 + z^2 + w^2$ for integers x , y , z , $w \geq 0$ and $(x, y, w, z) = 1$. It follows that $d^2 = A^2 + B^2 + C^2$ where $A = (x^2 + y^2 - z^2 - w^2)$, $B = 2 \cdot (x \cdot z + y \cdot w)$, and $C = 2 \cdot (x \cdot w - y \cdot z)$. It may be assumed that $A \geq 0$ and since d is odd, $A \neq 0$. If $(d, A) \neq 1$, then p divides $d + A = 2 \cdot (x^2 + y^2)$. Since $p = 3 \pmod{4}$, p therefore divides both x and y . But then, since $d = x^2 + y^2 + z^2 + w^2$, p also divides z and w which is impossible. Hence, $(d, A) = 1$.

Since

$$\left(\frac{A}{d}, \frac{B}{d}, \frac{C}{d}\right) + \left(\frac{A}{d}, -\frac{B}{d}, -\frac{C}{d}\right) = \left(\frac{2 \cdot A}{d}, 0, 0\right) \in \Sigma_0(\mathbb{Q}^3),$$

where $(2 \cdot A, d) = 1$, it follows that $(1/d, 0, 0) \in \Sigma_0(\mathbb{Q}^3)$ and therefore completes the assertion.

Example 3. $\Sigma_0(\mathbb{Q}^4)$ is the disjoint union of the sets $A = \mathbb{Z}_{(2)}^4$ and $B = \mathbb{Z}_{(2)}^4 + (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$.

Since A is a subgroup in \mathbb{Q}^4 and $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}) \notin A$, it follows that the sets are disjoint. Using Example 2, it is clear that all the vectors in the two sets belong to $\Sigma_0(\mathbb{Q}^4)$.

Let a, b, c, d , and e be integers for which $(a, b, c, d, e) = 1$ and $a^2 + b^2 + c^2 + d^2 = e^2$. If e is odd then $1/e \cdot (a, b, c, d) \in A$. If e is even, it follows by considering the equation modulo 8, that $e = 2 \pmod{4}$ and that a, b, c , and d are odd. Hence, $e = 2 \cdot f$ where f is odd and therefore $1/e \cdot (a, b, c, d) =$

$$\frac{1}{f} \cdot \left(\frac{a-f}{2}, \frac{b-f}{2}, \frac{c-f}{2}, \frac{d-f}{2} \right) + \left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \right),$$

where the first term of the sum belongs to A since the differences are all even integers. It is now clear that $\Sigma_0(\mathbb{Q}^4) \subset A \cup B$ which completes the assertion.

Theorem 5. \mathbb{Q}^4 has an additive 4-coloring.

Proof. From Example 3, it is clear that $\mathbb{Z}_{(2)}^4$ forms a subgroup of $\Sigma_0(\mathbb{Q}^4)$ and that $\Sigma_0(\mathbb{Q}^4) \cong \mathbb{Z}_{(2)}^4 \oplus \mathbb{Z}/(2)$. Let $s: \mathbb{Z}_{(2)}^4 \rightarrow \mathbb{Z}/(2)$ be the map which sums all the coordinates. Then $\epsilon \cdot s: \mathbb{Z}_{(2)}^4 \rightarrow \mathbb{Z}/(2)$ together with the identity map 1 on $\mathbb{Z}/(2)$, gives a group homomorphism

$$v = (\epsilon \cdot s, 1): \mathbb{Z}_{(2)}^4 \oplus \mathbb{Z}/(2) \rightarrow \mathbb{Z}/(2) \oplus \mathbb{Z}/(2).$$

Let $\xi \in \Sigma_0(\mathbb{Q}^4)$ with $\|\xi\| = 1$. According to Example 3, either $\xi = (a/e, b/e, c/e, d/e)$ with e odd or $\xi = (a/e, b/e, c/e, d/e) + (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ with e odd. In the former case, since also $a^2 + b^2 + c^2 + d^2 = e^2$, it follows that $a + b + c + d \neq 0 \pmod{2}$. Identifying ξ in the direct sum it follows that $v(\xi) \neq 0$.

In the latter case, ξ is identified as $(a/e, b/e, c/e, d/e) \oplus [1]$ and it follows again that $v(\xi) \neq 0$. Hence, v gives the required map. \square

It should be noted that since the points $(0, 0, 0, 0)$, $(1, 0, 0, 0)$, $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ and $(\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ form a tetrahedron in \mathbb{Q}^4 , the above shows that the chromatic number of \mathbb{Q}^4 is 4. This is known ([2], among others). Furthermore, since $\Sigma_0(\mathbb{Q}^d) = \mathbb{Q}^d$ for $d \geq 5$, it follows that \mathbb{Q}^d can have no additive k -coloring for any $k \geq 1$. For if $v: \mathbb{Q}^d \rightarrow G$ is an additive k -coloring, then $k \cdot v(1/k, 0, \dots, 0) = v(1) \neq 0$ and yet $k \cdot G = 0$ gives a contradiction.

The following theorem indicates that an arbitrary k -coloring as well as an additive k -coloring may be extended to a transcendental extension.

Theorem 6. *Let α be a real number transcendental over the field $F \subset \mathbb{R}$. Then F^d can be k -colored (has an additive k -coloring) iff $F(\alpha)^d$ can be k -colored (has an additive k -coloring).*

Proof. If $F(\alpha)^d$ can be k -colored, then surely so can F^d .

To show the converse, since α is transcendental over F , the field $F(\alpha)$ can be viewed as the rational function field $F(x)$ in the indeterminate x . Therefore, let

$$\xi(\alpha) = \left(\frac{p_1(\alpha)}{c(\alpha)}, \dots, \frac{p_d(\alpha)}{c(\alpha)} \right) \in F(\alpha)^d$$

where $c(x)$ and $p_j(x)$ for $j = 1, \dots, d$ are polynomials in $F[x]$. If $\|\xi(\alpha)\| = 1$, then it may be assumed that

$$\sum_{j=1}^d p_j(x)^2 = c(x)^2$$

and that these polynomials have no common factor.

Setting $x = 0$, one sees that if $c(0) = 0$, then $p_j(0) = 0$ for all j since all the polynomials have real coefficients. But since x would then be a common factor of all the polynomials, it follows that $c(0) \neq 0$. Hence,

$$\xi(0) = \left(\frac{p_1(0)}{c(0)}, \dots, \frac{p_d(0)}{c(0)} \right) \in F^d \quad \text{and still } \|\xi(0)\| = 1$$

because $\sum_{j=1}^d p_j(0)^2 = c(0)^2$.

In order to k -color $F(\alpha)^d$, it is clearly sufficient to k -color $\Sigma_0(F(\alpha)^d)$. Since the evaluation at 0 is linear, the above shows that if $\xi(\alpha) \in \Sigma_0(F(\alpha)^d)$, then $\xi(0) \in \Sigma_0(F^d)$. Assign to $\xi(\alpha)$ the color $\xi(0)$ which it receives from the assumed k -coloring of $\Sigma_0(F^d)$. To show that this gives a k -coloring of $\Sigma_0(F(\alpha)^d)$ it suffices to show that if $\xi_1(\alpha), \xi_2(\alpha) \in \Sigma_0(F(\alpha)^d)$ with $\xi_1(\alpha) - \xi_2(\alpha) = \xi(\alpha)$ and where $\|\xi(\alpha)\| = 1$, then $\xi_1(\alpha)$ and $\xi_2(\alpha)$ are assigned different colors. But then $\xi_1(0), \xi_2(0) \in \Sigma_0(F^d)$ with $\xi_1(0) - \xi_2(0) = \xi(0)$ and where $\|\xi(0)\| = 1$. But since F^d can be k -colored, $\xi_1(0)$ and $\xi_2(0)$ must receive different colors and hence also $\xi_1(\alpha)$ and $\xi_2(\alpha)$.

To show the assertion for an additive k -coloring, it is sufficient to show that the given map $v: \Sigma_0(F^d) \rightarrow G$ may be extended to a map $\bar{v}: \Sigma_0(F(\alpha)^d) \rightarrow G$. Using the notation above, if $\xi(\alpha) \in \Sigma_0(F(\alpha)^d)$, then define $\bar{v}(\xi(\alpha)) = v(\xi(0))$. Since the evaluation at 0 is a mapping of groups, so is \bar{v} . As above, if $\|\xi(\alpha)\| = 1$, then $\|\xi(0)\| = 1$ and it follows that $\bar{v}(\xi(\alpha)) = v(\xi(0)) \neq 0$. \square

Remark 1. Given a finitely generated extension $E \subset \mathbb{R}$ over \mathbb{Q} , it may be realized as a purely transcendental extension of some finite algebraic extension F over \mathbb{Q} . It follows from the above theorem that E^d can be k -colored iff F^d can be k -colored. Given a finite number of points in \mathbb{R}^d , one may adjoin the coordinates of these points to form a finitely generated extension E over \mathbb{Q} . Using the result

of de Bruijn and Erdős as stated in the introduction, it follows that the question of the k -colorability of an arbitrary extension of \mathbb{Q} may be reduced to that of the k -colorability of a finite algebraic extension of \mathbb{Q} .

Remark 2. Again using the result of de Bruijn and Erdős and the above theorem, it can be seen that there exist fields F_0 so that F_0^d is maximal in respect to being k -colored provided that \mathbb{Q}^d is k -colorable. Because of the above theorem, \mathbb{R} must also be algebraic over F_0 .

3. Real quadratic extensions of \mathbb{Q}

Any real quadratic extension of \mathbb{Q} may be realized as $\mathbb{Q}(\sqrt{N})$ where N is a square free positive integer. In this section, the additive k -colorability of the plane $\mathbb{Q}(\sqrt{N})^2$ will be investigated. In what follows, Σ_0 will always denote $\Sigma_0(\mathbb{Q}\sqrt{N})^2$ and for an integer n , $[n]$ will denote the equivalence class of the integer n modulo 2.

The first theorem can be used as a test of whether a plane is 2-colorable.

Theorem 7. Suppose that the plane F^2 can be 2-colored and hence has a ring homomorphism $v: \Sigma_0(F^2) \rightarrow \mathbb{Z}/(2)$. Let $p(x) = a_n \cdot x^n + \cdots + a_0$ be a polynomial so that $a_k \in \Sigma_0(F^2)$ for each $k = 1, \dots, n$. If $p(\xi) = 0$ for some $\xi \in \Sigma_0(F^2)$ with $v(\xi) = [1]$, then $\sum_{k=0}^n v(a_k) = [0]$.

In particular, if $p(x) = a_n \cdot x^n + \cdots + a_0$ is a polynomial with integral coefficients for which $p(\xi) = 0$ for some $\xi \in \mathbb{Q}(i)$ with $\|\xi\| = 1$, then $a_n + \cdots + a_0 = 0 \pmod{2}$.

Proof. Since $\xi \in \Sigma_0(F^2)$ so is ξ^k for $k = 0, \dots, n$ and hence $p(\xi) \in \Sigma_0(F^2)$. Clearly, $v(\xi^k) = v(\xi)$ for all k since v is a homomorphism. Applying v to the equation $0 = p(\xi) = a_n \cdot \xi^n + \cdots + a_0$ now gives the first result.

If the a_k 's are integers then $a_k \in \Sigma_0(F^2)$ and $v(a_k) = [n]$. By assumption $v(\xi) = [1]$ and the second result follows from the first. \square

A recent and independent result of Johnson [8] shows in more generality the “if” portion of the following theorem. He shows that the points in $\mathbb{Q}(\sqrt{N})^2$ for $N \not\equiv 3 \pmod{4}$ may be partitioned into sets A and B which fail to realize not only the distance 1 but also the distances $\sqrt{p/q}$ where p and q are positive odd integers.

In what follows, recall that for an integer n , $n^2 = 1 \pmod{8}$ iff n is odd, $n^2 = 4 \pmod{8}$ iff $n = 2 \pmod{4}$, and that $n^2 = 0 \pmod{8}$ iff $n = 0 \pmod{4}$.

Theorem 8. Let N be a square free positive integer. Then the plane $\mathbb{Q}(\sqrt{N})^2$ can be 2-colored iff $N \not\equiv 3 \pmod{4}$.

Proof. If N is a positive integer for which $N \equiv 3 \pmod{4}$, then $N + 1 = 4 \cdot a$ for some integer $a > 0$. It follows that

$$N = \left[\frac{n+1}{2} \right]^2 - \left[\frac{N-1}{2} \right]^2 = 4 \cdot a^2 - b^2,$$

where $b = (N - 1)/2$ must be odd. But then

$$\xi = \frac{b + \sqrt{N} \cdot i}{2 \cdot a}$$

satisfies $\|\xi\| = 1$ and is a solution of $a \cdot x^2 - b \cdot x + a = 0$. Since $2 \cdot a - b \equiv 1 \pmod{2}$, the plane $\mathbb{Q}(\sqrt{N})^2$ where $N \equiv 3 \pmod{4}$ cannot be 2-colored because otherwise this would contradict Theorem 7.

In order to argue the converse, it suffices by Theorem 1 to find an additive map $v: \Sigma_0 \rightarrow \mathbb{Z}/(2)$ for which $v(\xi) = [1]$ whenever $\|\xi\| = 1$. Let $\mathbb{Z}_{(2)}$, as before, be the subring of those fractions of \mathbb{Q} with odd denominators and ϵ the evaluation $\epsilon: \mathbb{Z}_{(2)} \rightarrow \mathbb{Z}/(2)$.

Any $\xi \in \mathbb{Q}(\sqrt{N}, i)$ may be written as

$$\xi = \frac{(a_1 + b_1 \cdot i) + (a_2 + b_2 \cdot i)\sqrt{N}}{c},$$

where c, a_j , and b_j for $j = 1, 2$ are integers and for which it may be assumed that $(a_1, b_1, a_2, b_2, c) = 1$. If $\|\xi\| = 1$, then $\xi \cdot \bar{\xi} = 1$. Performing the multiplication and noticing that $\sqrt{N} \notin \mathbb{Q}$, one obtains the equations

$$a_1^2 + b_1^2 + N \cdot (a_2^2 + b_2^2) = c^2 \quad \text{and} \quad (1)$$

$$a_1 \cdot a_2 + b_1 \cdot b_2 = 0. \quad (2)$$

The case $N \equiv 1 \pmod{4}$ will be argued first and the various four exhaustive possibilities for $[a_1]$ and $[b_1]$ considered.

If $[a_1] = [b_1] = [1]$, equation (1) taken modulo 8 gives $c^2 = a_1^2 + b_1^2 + N \cdot (a_2^2 + b_2^2) \equiv 2 + N \cdot (a_2^2 + b_2^2) \pmod{8}$. Since $N \equiv 1 \pmod{4}$, it can be seen from this that necessarily $a_2^2 = b_2^2 \equiv 1 \pmod{8}$ and that $c^2 \equiv 4 \pmod{8}$. Hence, $[a_1] = [b_1] = [a_2] = [b_2] = [1]$ and $c/2$ is an odd integer. Since $a_1 \cdot a_2 + b_1 \cdot b_2 = 0$ and since all the integers are assumed to be odd, it follows that

$$\begin{aligned} (a_1 + b_1 + a_2 + b_2)^2 &= a_1^2 + b_1^2 + a_2^2 + b_2^2 + 2 \cdot (a_1 + a_2)(b_1 + b_2) \\ &\equiv 4 \pmod{8}. \end{aligned}$$

Hence, $(a_1 + b_1 + a_2 + b_2)/2$ is also an odd integer and shows

$$\frac{a_1 + b_1 + a_2 + b_2}{c} \in \mathbb{Z}_{(2)} \quad \text{and} \quad \epsilon\left(\frac{a_1 + b_1 + a_2 + b_2}{c}\right) = [1]. \quad (3)$$

But the same conclusion (3) holds true in the remaining possibilities of $[a_1]$ and $[b_1]$. For if $[a_1 + b_1] = [1]$, then equation (1) taken modulo 4 shows that it has to

be that $[a_2] = [b_2] = [0]$ and that $[c] = [1]$. Hence, c is odd and $[a_1 + b_1 + a_2 + b_2] = [1]$.

Finally, if $[a_1] = [b_1] = [0]$ then Equation (1) gives $c^2 = a_2^2 + b_2^2 \pmod{4}$. If $[c] = 0$, then this would mean $[a_2] = [b_2] = [0]$ contradicting the fact that $(a_1, b_1, a_2, b_2, c) = 1$. Hence, it must be that $[c] = [1]$ and that $[a_2 + b_2] = [1]$. Again, therefore, conclusion (3) holds true.

Using the identification $\mathbb{Q}(\sqrt{N}, i) \cong \mathbb{Q}^4$ along with the linear functional s which sums the coordinates of \mathbb{Q}^4 , one has an additive map

$$s: \mathbb{Q}(\sqrt{N}, i) \rightarrow \mathbb{Q} \quad \text{where } s(\xi) = \frac{a_1 + b_1 + a_2 + b_2}{c}.$$

The results from (3) show that $s(\Sigma_0) \subset \mathbb{Z}_{(2)}$ and that $\epsilon(s(\xi)) = [1]$ if $\|\xi\| = 1$. Hence,

$$v = \epsilon \cdot s: \sum_0 \rightarrow \mathbb{Z}/(2)$$

is the desired group map in the case that $N \equiv 1 \pmod{4}$.

For the case $N \equiv 2 \pmod{4}$, one notes that if $[a_1] = [b_1] = [1]$, then equation (2) shows that $[a_2] = [b_2]$. It follows by considering equation (1) modulo 4, that no ξ for which $\|\xi\| = 1$ and $[a_1] = [b_1] = [1]$ can exist in this case.

If $[a_1 + b_1] = [1]$, then equation (1) shows that $[c] = [1]$ and $[a_2] = [b_2] = [0]$. Hence,

$$\frac{a_1 + b_1 + 2 \cdot b_2}{c} \in \mathbb{Z}_{(2)} \quad \text{and} \quad \epsilon\left(\frac{a_1 + b_1 + 2 \cdot b_2}{c}\right) = [1]. \quad (4)$$

But the same conclusion as in (4) holds for the remaining case of when $[a_1] = [b_1] = [0]$. For then Equation (1) implies that $[c] = [0]$ and that $[a_2] = [b_2]$. It cannot be that $[a_2] = [b_2] = [0]$ because $(a_1, b_1, a_2, b_2, c) = 1$. Hence, $[a_2] = [b_2] = [1]$. Observe, therefore, that since $a_1 \cdot a_2 + b_1 \cdot b_2 = 0$, a_1 and b_1 contain the same power of 2 in their factorization and consequently $a_1 + b_1 \equiv 0 \pmod{4}$ or equivalently, $(a_1 + b_1)/2$ is an even integer. Furthermore, since also $a_2^2 + b_2^2 \equiv 2 \pmod{8}$, equation (1) yields because $N \equiv 2 \pmod{4}$ that $c^2 \equiv 0 + N \cdot 2 \equiv 4 \pmod{8}$. Hence, $c/2$ is an odd integer. Since $[b_2] = [1]$, the results stated in (4) hold in this case also.

Again identifying $\mathbb{Q}(\sqrt{N}, i)$ with \mathbb{Q}^4 , one has the map

$$t: \mathbb{Q}(\sqrt{N}, i) \rightarrow \mathbb{Q} \quad \text{where}$$

$$t(\xi) = \frac{a_1 + b_1 + 2 \cdot b_2}{c}.$$

The results in (4) show therefore, that $t(\Sigma_0) \subset \mathbb{Z}_{(2)}$ and that $\epsilon \cdot t(\xi) = [1]$ if $\|\xi\| = 1$. Hence, $v = \epsilon \cdot t: \Sigma_0 \rightarrow \mathbb{Z}/(2)$ is the required map in the case $N \equiv 2 \pmod{4}$. \square

The equations appearing in the above theorem can be utilized to demonstrate some additional k -colorings of quadratic extensions.

Theorem 9. *Suppose N is a square free positive integer for which $N \not\equiv 2 \pmod{3}$. Then the plane $\mathbb{Q}(\sqrt{N})^2$ has an additive 3-coloring.*

It follows that the chromatic number of the plane $\mathbb{Q}(\sqrt{N})^2$ is 3 when $N = 3$ or $7 \pmod{12}$.

Proof. As in the above theorem, if

$$\xi = \frac{(a_1 + b_1 \cdot i) + (a_2 + b_2 \cdot i)\sqrt{N}}{c} \in \mathbb{Q}(\sqrt{N}, i)$$

with $\|\xi\| = 1$, then

$$a_1^2 + b_1^2 + N \cdot (a_2^2 + b_2^2) = c^2 \quad \text{and} \quad (1)$$

$$a_1 \cdot a_2 + b_1 \cdot b_2 = 0 \quad (2)$$

where the integers a_1, b_1, a_2, b_2 and c have no common non-trivial divisor.

Let $\mathbb{Z}_{(3)}$ be the local ring of fractions whose denominator is not divisible by 3 and let $\delta: \mathbb{Z}_{(3)} \rightarrow \mathbb{Z}/(3)$ be the evaluation at the maximal ideal (3) . That is, for $a/b \in \mathbb{Z}_{(3)}$ with $b \not\equiv 0 \pmod{3}$, $\delta(a/b) = \bar{a} \cdot (\bar{b})^{-1}$ where \bar{a} and \bar{b} are the equivalence classes of the integers a and b respectively in the ring $\mathbb{Z}/(3)$ of the integers modulo 3.

If $N \equiv 0 \pmod{3}$, note that equation (1) when taken modulo 3 shows that $\bar{c} \neq \bar{0}$. For if $\bar{c} = \bar{0}$, then since $\bar{N} = \bar{0}$, $\bar{a}_1^2 + \bar{b}_1^2 = \bar{0}$ which implies that a_1 and b_1 are divisible by 3. It follows that 3^2 divides $N \cdot (a_2^2 + b_2^2)$ and that therefore $\bar{a}_2^2 + \bar{b}_2^2 = \bar{0}$ since N is square free. Hence, a_2 and b_2 are also divisible by 3 which contradicts the assumption that $(a_1, b_1, a_2, b_2, c) = 1$.

Hence, $\bar{c} \neq \bar{0}$ and therefore $\bar{c}^2 = \bar{1}$. It follows again by equation (1) that $\bar{a}_1^2 + \bar{b}_1^2 = \bar{1}$ and that therefore $\bar{a}_1 + \bar{b}_1 \neq \bar{0}$.

Identifying $\mathbb{Q}(\sqrt{N}, i)$ with \mathbb{Q}^4 and setting

$$t\left(\frac{a_1 + b_1 i + (a_2 + b_2 i)\sqrt{N}}{c}\right) = \frac{a_1 + b_1}{c},$$

the above shows that if $N \equiv 0 \pmod{3}$ then $v = \delta \cdot t: \Sigma_0 \rightarrow \mathbb{Z}/(3)$ is the desired additive map with $v(\xi) \neq \bar{0}$ whenever $\|\xi\| = 1$.

If $N \equiv 1 \pmod{3}$, note that Equation (1) taken modulo 3 shows that

$$\bar{a}_1^2 + \bar{b}_1^2 + \bar{a}_2^2 + \bar{b}_2^2 = \bar{c}^2.$$

If $\bar{c} = \bar{0}$, then at least one of $\bar{a}_1, \bar{b}_1, \bar{a}_2$, or \bar{b}_2 must equal $\bar{0}$. But then Equation (2) set to modulo 3 shows that another must equal $\bar{0}$ also. It follows that a_1, b_1, a_2, b_2 and c are all divisible by 3 contradicting the assumption that these have no nontrivial divisor. Hence, $\bar{c} \neq \bar{0}$ and $\bar{c}^2 = \bar{1}$.

By equation (2) again,

$$(a_1 + b_1 + a_2 + b_2)^2 = a_1^2 + b_1^2 + a_2^2 + b_2^2 + 2 \cdot (a_1 + a_2)(b_1 + b_2).$$

If $\bar{a}_1 + \bar{b}_1 + \bar{a}_2 + \bar{b}_2 = \bar{0}$ then also $(\bar{a}_1 + \bar{a}_2) = -(\bar{b}_1 + \bar{b}_2)$ and hence $\bar{0} = \bar{1} + (-\bar{2})(\bar{a}_1 + \bar{a}_2)^2$. But this is impossible and therefore it must be that $\bar{a}_1 + \bar{b}_1 + \bar{a}_2 + \bar{b}_2 \neq \bar{0}$.

It follows in the case $N = 1 \pmod{3}$ that if

$$s\left(\frac{a_1, b_1, a_2, b_2}{c}\right) = \frac{a_1 + b_1 + a_2 + b_2}{c}, \quad \text{then}$$

$v = \delta \cdot s : \Sigma_0 \rightarrow \mathbb{Z}/(3)$ is the desired map with $v(\xi) \neq \bar{0}$.

The second part of the theorem now follows from Theorem 8 since to say that $N = 3$ or $7 \pmod{12}$ is equivalent to asserting that $N = 3 \pmod{4}$ and $N \neq 2 \pmod{3}$. \square

The existence of an additive k -coloring on quadratic extensions for those cases not considered by the previous two theorems is covered in the following.

Theorem 10. Suppose N is a positive square free integer. Then the plane $\mathbb{Q}(\sqrt{N})^2$

- i. has an additive 4-coloring if $N = 3 \pmod{8}$.
- ii. has no additive k -coloring for $1 \leq k \leq 6$ if

$$N = -1 \pmod{24}.$$

Proof. Equations (1) and (2) referred to in the following are those as in the previous theorems.

To prove (i), note that if $c = 1 \pmod{2}$, then equation (1) taken modulo 4 shows that since $N = 3 \pmod{4}$, either $a_1 + b_1 = 1$ and $a_2 = b_2 = 0 \pmod{2}$ or that $a_1 = b_1 = 1 \pmod{2}$ and $a_2 + b_2 = 1 \pmod{2}$. But the latter situation cannot occur because of equation (2). Hence, if $\epsilon : \mathbb{Z}_{(2)} \rightarrow \mathbb{Z}/(4)$ is the map which evaluates at the square of the maximal ideal, one has in this case:

$$\frac{2(a_1 + b_1)}{c} \in \mathbb{Z}_{(2)} \quad \text{and} \quad \epsilon\left(\frac{2(a_1 + b_1)}{c}\right) \neq 0. \quad (3)$$

If $c = 0 \pmod{2}$, then not all of the integers a_1, b_1, a_2 , and b_2 can equal $0 \pmod{2}$. It follows by equation (1) taken modulo 4, that either $a_1 + b_1 = a_2 + b_2 = 1 \pmod{2}$ or that $a_1 = b_1 = a_2 = b_2 = 1 \pmod{2}$. If the latter, then because of equation (1) taken modulo 8 and since $N = 3 \pmod{8}$, it follows that $c^2 = 0 \pmod{8}$ and therefore $c = 4 \cdot m$ for some integer m . Since in this case $a_1 \neq 0$, one has

$$(a_2, b_2) = \frac{b_2}{a_1}(-b_1, a_1)$$

because of Equation (2). Substituting this into Equation (1) yields

$$(a_1^2 + b_2^2 N)(a_1^2 + b_1^2) = a_1^2 \cdot c^2.$$

But since $a_1^2 + b_1^2 N = 4 \pmod{8}$ and since $c^2 = 16 \cdot m^2$, it follows that $a_1^2 + b_1^2 = 0 \pmod{4}$ which is impossible since a_1 and b_1 are odd integers.

Hence, it may be assumed that if $c = 0 \pmod{2}$, then $a_1 + b_1 = a_2 + b_2 = 1 \pmod{2}$. It follows from this and Equation (2) that $a_1^2 + b_1^2 = a_2^2 + b_2^2 = 1$ or $5 \pmod{8}$. But then Equation (1) taken modulo 8 shows that since $N = 3 \pmod{8}$, $c^2 = 4 \pmod{8}$ and hence $c = 2 \cdot m$ where m is an odd integer. It follows again that

$$\frac{2(a_1 + b_1)}{2 \cdot m} = \frac{(a_1 + b_1)}{m} \in \mathbb{Z}_{(2)} \quad \text{and} \quad \epsilon\left(\frac{2(a_1 + b_1)}{c}\right) \neq 0.$$

Identifying $\mathbb{Q}(\sqrt{N}, i)$ with \mathbb{Q}^4 and setting

$$t\left(\frac{a_1, b_1, a_2, b_2}{c}\right) = \frac{2 \cdot (a_1 + b_1)}{c},$$

then the above shows that $t: \Sigma_0 \rightarrow \mathbb{Z}_{(2)}$ and that $v = \epsilon \cdot t: \Sigma_0 \rightarrow \mathbb{Z}/(4)$ is the desired map for which $v(\xi) \neq 0$ if $\|\xi\| = 1$.

To prove (ii), note that since $N = a^2 - b^2$ where $a = (N+1)/2$ and $b = (N-1)/2$, it follows from the assumptions on N that $a = 3 \cdot 4 \cdot m$ for some integer m and that $b = 1 \pmod{2}$ and $b = 2 \pmod{3}$.

Since for $\xi = (b + \sqrt{N} \cdot i)/a$, $\|\xi\| = 1$, it follows that $\xi + \bar{\xi} = (2 \cdot b)/a = b/(2 \cdot 3 \cdot m) \in \Sigma_0$. But since $(2, b) = 1 = (3, b)$ it must be that $\frac{1}{2}$ and $\frac{1}{3}$ are in Σ_0 as is $\frac{1}{5} \in \Sigma_0(\mathbb{Q}^2) \subset \Sigma_0$. Since Σ_0 is a ring, $1/k \in \Sigma_0$ for $1 \leq k \leq 6$.

As noted before, if v is an additive k -coloring $v: \Sigma_0 \rightarrow G$ to some group G of order k , then $k \cdot v(1/k) = v(1) \neq 0$. But $k \cdot G = 0$ and hence no such map can exist. \square

Example 4. The integer $167 = 3 \cdot 7 \cdot 2^3 - 1$ is the first one which satisfies the hypothesis of part (ii) of the theorem and also equals $6 \pmod{7}$. Therefore, if

$$\xi = \frac{83 + \sqrt{167} \cdot i}{3 \cdot 7 \cdot 2^2}$$

then $\|\xi\| = 1$. Hence, $\frac{1}{2}$, $\frac{1}{3}$, and $\frac{1}{7}$ are in Σ_0 .

As in part (ii) of the theorem, this shows that $\mathbb{Q}(\sqrt{167})^2$ has no additive k -coloring for $k < 11$. Hence, even though the plane formed by the quadratic extension $\mathbb{Q}(\sqrt{167})$ can be 7-colored since \mathbb{R}^2 can, it has no additive 7-coloring.

Acknowledgments

I wish to thank Peter D. Johnson, Jr. for furnishing the appropriate papers and references in this area, his comments, and also pointing out a flaw in the original manuscript. Thanks also to the referee for the very careful reading of this paper and to the Department of Mathematics at the University of Illinois at Urbana for its hospitality during the writing of this paper.

References

- [1] N.G. de Bruijn and P. Erdős, A colour problem for infinite graphs and a problem in the theory of relations, *Nederl. Akad. Wetensch. Indag. Math.*, 13 (1951) 371–373.
- [2] M. Benda and M. Perles, Colorings of metric spaces (preprint).
- [3] K.B. Chilakamarri, Unit-distance graphs in rational n -spaces, *Discrete Mathematics* 69 (1988) 213–218.
- [4] E. Dickson, *History of the Theory of Numbers*, vol. II (Chelsea, New York, 1971).
- [5] P. Erdős, Some new problems and results in graph theory and other branches of combinatorial mathematics, *Combinatorics and Graph Theory*, Lecture Notes in Mathematics, no. 885 (1981).
- [6] Hadwiger, Debrunner and Klee, *Combinatorial Geometry in the Plane* (Holt, Rinehart and Winston, New York, 1964).
- [7] P.D. Johnson, Jr., Coloring abelian groups, *Discrete Mathematics* 40 (1982) 219–223.
- [8] P.D. Johnson, Jr., Two-colorings of real quadratic extensions of \mathbb{Q}^2 that forbid many distances, *Congressus Numerantium, the Proceedings of the Eighteenth Southeastern Conference on Combinatorics, Graph Theory and Computing* (to appear).
- [9] D.R. Woodall, Distances realized by sets covering the plane, *J. Comb. Theory, Ser. A*, 14 (1973) 187–200.